

# 조직생검된 사람 정상 코점막 상피세포와 배양된 코점막 상피세포에서 점액과 리소자임 단백질의 정량 및 유전자 발현 비교

윤주현 · 김경수 · 김성식 · 김재원 · 이정권 · 박인용

## Comparison of Mucin and Lysozyme Expression between Human in vivo Nasal Epithelial Cells and Cultured Nasal Epithelial Cells

Joo-Heon Yoon, MD, Kyung-Su Kim, MD, Sung-Shik Kim, MD, Jae Won Kim, MD, Jeung-Gweon Lee, MD and In Yong Park, MD

Department of Otorhinolaryngology Yonsei University College of Medicine, Seoul, Korea

### ABSTRACT

**Background and Objectives :** The purpose of this study was to compare the expression of mucin and lysozyme in passage-2 normal human nasal epithelial (NHNE) cells with those in human in vivo nasal epithelium and human tracheal RNA. **Materials and Methods :** Cell lysates and total RNA from passage-2 NHNE cells, and human in vivo nasal epithelial cells were obtained. The amount of mucin and lysozyme protein was measured by immunoblotting, and qualitative RT-PCR was done to investigate the expression of mucin mRNAs and lysozyme mRNA. **Results :** Passage-2 NHNE cells contained 16% of mucin and 76% of lysozyme when compared to the amount of intracellular mucin and lysozyme of normal in vivo nasal epithelial cells. MUC4, MUC5AC, MUC7, MUC8 and lysozyme mRNAs were expressed in passage-2 NHNE cells. However, MUC2 and MUC5B mRNAs were not expressed. **Conclusion :** Passage-2 NHNE cells contain enough amount of mucin and lysozyme protein and express most mRNAs of secretory genes which are known to be expressed in the human airway. Thus, we find passage-2 NHNE cells to be suitable for conducting studies on secretions in the human upper airway. (Korean J Otolaryngol 1999;42:317-21)

**KEY WORDS :** Normal human nasal epithelial cells · Mucin · Lysozyme · Protein · mRNA..

retinoic acid(RA)가

1) 2) 3)

passage - 2 air - liquid interface(ALI)

가

4)

(passage - 2)

가 (lower

airway)

: 1998 10 15 / : 1999 1 30

: , 120 - 752 134

: (02) 361 - 8470 · : (02) 393 - 0580

E - mail : jhyoon@yumc.yonsei.ac.kr

ge - 2 , , passa - North Carolina, NC, USA) (Sigma, St. Louis, MO) MO)

H6C5(1 : 1,000, a generous gift from Dr. Davis CW)

MUC1, MUC3 MUC6 9 (1 : 1,000, Dako, Capintera, CA)

mRNA mRNA 6

Passage - 2 horse - radish peroxidase conjugated goat anti - mouse IgG anti - rabbit IgG che - miluminescence(ECL kit, Amersham, Buckinghamshire, UK) . Linear regression analysis standard curve

(Transwell - clear, Costar Corp., Cambri - dge, MA) 3.0 mg/ml 1 (Collaborative Res., New Bedford, MA) ammonium hydroxide Western blot , 10<sup>5</sup>

(normal human nasal epithelial(NHNE) hemocytometer cells, passage - 2) 가 가 3 ± 4) 9 Student's t - test ALI 9

37 0C, 5% CO<sub>2</sub> PCR mRNA RT -

Transwell - clear membrane MUC1 in situ

16 3 wells cell lysate hybridization MUC3 MUC6 . Oligonucleotide RNA 20 30 10 primers human

scraping 37 1 MUC2(Genbank accession #L21998, 5' primer : TG - CCTGGCCCTGTCTTTG ; 3' primer : CAGCTCCA - GCATGAGTGC) ; human MUC4(Genbank accession #AJ000281, 5' primer : TTCTAAGAACCACCAG - ACTCAGAGC ; 3' primer : GAGACACACCTGGAG - AGAATGAGC) ; human MUC5AC(Genbank accession #U06711, 5' primer : TCCGGCTCATCTTCTTC - C ; 3' primer : ACTTGGGCACTGGTGCTG) ; humanM - UC5B(Genbank accession #Z72496, 5' primer : ACT CCAGAGACTGTCCACAC ; 3' primer : TACC - ACT GGTCTGTGTGCTA) ; human MUC7(Genbank acce - ssion #L132283, 5' primer : CCACACCTAA - TTC TTCCC ; 3' primer : CTATTGCTCCACCATG - TC) ; human MUC8(Genbank accession #U14383, 5' pri - mer : ACAGGGTTTCTCCTCATTG ; 3' primer : CGT TTATTCCAGCACTGTTC) ; human lysozyme (Gen - bank accession #J03801, 5' primer : CTCTCAT - TG

trypsin/EDTA RNA

Tri - reagent cell lysate

total RNA, Clontech, Palo Alto, CA, USA) RNA(human trachea

24 immunoblot assay Gray 5) RA


(a generous gift from Dr. Davis CW, University of

TTCTGGGGC ; 3' primer : ACGGACA ACCCTC - TT  
TGC) . cDNA MUC2  
440 bp, MUC4 466 bp, MUC5AC 680 bp,  
MUC5B 338 bp, MUC7 209 bp, MUC8 239 bp,  
350 bp . Control gene 2  
micro - globulin 350 bp . RT - PCR  
Perkin Elmer Cetus DNA Thermal Cycler  
primer kinetic study  
MgCl<sub>2</sub> , MUC4, MUC5B, MUC5AC,  
MUC7, MUC8, 2M 1.5 mM  
MUC2 2.5 mM , MUC2, MUC - 5AC  
2M annealing temperature 60 , MUC4, MU C5B,  
MUC7, MUC8 55 C .  
linear range MUC2 29 35  
cycles, MUC5AC 32 40 cycles, MUC5B 23 29  
cycles, MUC7 32 40 cycles, MUC8 32 38  
cycles . PCR genomic DNA  
RT reaction RT enzyme  
PCR band가  
DNA sequencing(dsDNA Cycle Sequ -  
encing System, Gibco - BRL) .

**Table 1.** Comparison of mucin and lysozyme protein levels between pooled normal human in vivo nasal epithelial cells and fully-differentiated passage-2 normal human nasal epithelial cells

	Mucin	Lysozyme
Pooled normal in vivo nasal epithelial cells	814.9	10.0
Fully-differentiated passage-2 nasal epithelial cells	131.4 ± 9.1	7.6 ± 2.3

value : µg/10<sup>6</sup>cells



**Fig. 1.** Comparison of mucin mRNA levels and lysozyme mRNA levels in normal human nasal epithelial cells from in vivo tissue (A), fully-differentiated passage-2 NHNE cells (B) and human tracheal RNA (C) by RT-PCR. The levels of MUC2, MUC5B, MUC8, and lysozyme mRNA decreased in fully-differentiated passage-2 NHNE cells compared to normal human nasal epithelial cells from in vivo tissue. However, MUC5AC mRNA levels were not altered. MUC4 mRNA levels were strongly expressed in passage-2 NHNE cells. MUC7 mRNA levels were strongly expressed in human tracheal RNA compared to human nasal epithelial cells whether they were cultured or not. MUC8 mRNA levels were strongly expressed in normal human nasal epithelial cells from in vivo tissue compared to fully-differentiated passage-2 NHNE cells and human tracheal RNA. Control gene, 2M, was not altered.

10<sup>6</sup> 814.9 µg ,  
10<sup>6</sup> 131.4 ± 9.1 µg  
가 16.1%  
10<sup>6</sup> 10 µg ,  
10<sup>6</sup> 7.6 ± 2.3 µg  
가 76%  
(Table 1).

MUC8

RNA 가 가 (Fig. 1).  
MUC2  
MUC5B가  
MUC4  
MUC5AC MUC7  
passage - 2  
RA가

가 RA 가 RNA , in situ hybridization

<sup>5)</sup> 가 <sup>8)9)</sup> MUC2 MUC5AC

가 가 가

MUC8 MUC5B MUC7 MUC5B

Yoon <sup>4)</sup> , MUC7 <sup>10)</sup> MUC5B , MUC7 MUC4

Yoon 가 MUC4가

$10^6$  814.9  $\mu$ g 131.4  $\mu$ g 가 가 passage - 2 , MUC2 MUC5B가

가 , Gray passage - 2 가

<sup>5)</sup> passage - 2  $10^6$  100 400  $\mu$ g passage -

2

PCR RT - Northern , 가

blotting RT - PCR band 가

Northern blot smear cassette

<sup>6)</sup> mRNA 2 <sup>11)</sup>

가 가

MUC1 MUC8 9 가

1 passage - 2

MUC3 MUC1

MUC6, RNA RNA

## REFERENCES

- 1) Adler KB, Cheng PW, Kim KC. *Characterization of guinea pig tracheal epithelial cells maintained in biphasic organotypic culture: Cellular composition and biochemical analysis of released glycoproteins.* *Am J Respir Cell Mol Biol* 1990;2:145-54.
- 2) Kaartinen L, Nettesheim P, Adler KB, Randel SH. *Rat tracheal epithelial cell differentiation in vitro.* *In Vitro Cell Dev Biol* 1993;29A:481-92.
- 3) Niles R, Kim KC, Hyman B, Christensen T, Wasano K, Brody J. *Characterization of extended primary and secondary cultures of hamster tracheal epithelial cells.* *In Vitro Cell Dev Biol* 1988;24:457-63.
- 4) Yoon JH, Kim KS, Kim SS, Lee JG, Park IY. *Secretory Differentiation of Serially Passaged Normal Human Nasal Epithelial (NHNE) Cells.* *Kor J Otolaryngol* 1998;41:1008-14.
- 5) Gray T, Guzman K, Davis CW, Abdullah L, Nettesheim P. *Mucociliary differentiation of serially passaged normal human tracheobronchial epithelial cells.* *Am J Respir Cell Mol Biol* 1996;14:104-12.
- 6) Guzman K, Gray T, Yoon JH, Nettesheim P. *Quantitation of mucin RNA by PCR reveals induction of both MUC2 and MUC5AC mRNA levels by retinoids.* *Am J Physiol* 1996;271 (Lung Cell Mol Physiol 15):L1023-L1028.
- 7) Yoon JH, Gray T, Guzman K, Koo JS, Nettesheim P. *Regulation of the secretory phenotype of human airway epithelium by retinoic acid, triiodothyronine, and extracellular matrix.* *Am J Respir Cell Mol Biol* 1997;16:724-31.
- 8) Aust MR, Madson CS, Jennings A, Kasperbauer JL, Gendler SJ. *Mucin mRNA expression in normal and vasomotor inferior turbinates.* *Am J Rhinol* 1997;301:293-302.
- 9) Reid CJ, Gould S, Harris A. *Developmental expression of mucin genes in the human respiratory tract.* *Am J Respir Cell Mol Biol* 1997;17:592-8.
- 10) Sharma P, Dudus L, Nielson PA, Clausen H, Yankaskas JR, Hollingsworth MA, et al. *MUC5B and MUC7 are differentially expressed in mucous and serous cells of submucosal glands in human bronchial airways.* *Am J Respir Cell Mol Biol* 1998;19:30-7.
- 11) Yoon JH, Koo JS, Norford D, Guzman K, Gray T, Nettesheim P. *Lysozyme expression during metaplastic squamous differentiation of retinoic acid-deficient human tracheobronchial epithelial cells.* *Am J Respir Cell Mol Biol* (in press).